



Contents lists available at ScienceDirect

Sleep Health

Journal of the National Sleep Foundation

journal homepage: sleephealthjournal.org

Poor sleep quality and physical performance in older adults

Hayley J. Denison, PhD^{a,b,*}, Karen A. Jameson, MSc^a, Avan A. Sayer, MBBS, PhD, FRCP^{a,c,d},
Harnish P. Patel, BM, PhD, FRCP^{a,e,f}, Mark H. Edwards, MD, PhD^{a,g},
Teresa Arora, PhD, CSci, CPsychol, AFBPs^h,
Elaine M. Dennison, MB, BChir, MA, FRCP, MSc, PhD^a,
Cyrus Cooper, MA, DM, FRCP, FFPH, FMedSci^{a,f}, Janis Baird, MB, BCh, PhD, FFPH^{a,f}

^a Medical Research Council Lifecourse Epidemiology Unit, University of Southampton, Southampton, UK

^b Centre for Public Health Research, Massey University, PO Box 756, Wellington 6140, New Zealand

^c NIHR Newcastle Biomedical Research Centre, Newcastle upon Tyne Hospitals NHS Foundation Trust and Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne, UK

^d AGE Research Group, Newcastle University Translational and Clinical Research Institute, Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne, UK

^e Academic Geriatric Medicine, University of Southampton, Southampton, UK

^f NIHR Southampton Biomedical Research Centre, University of Southampton and University Hospital Southampton NHS Foundation Trust, Southampton, UK

^g Portsmouth Hospitals NHS Trust, Portsmouth, UK

^h Zayed University, College of Natural and Health Sciences, Abu Dhabi, United Arab Emirates

ARTICLE INFO

Keywords:

Sleep
Physical performance
Grip strength
Sarcopenia
Muscle
Aging

ABSTRACT

Objectives: This study aimed to examine the association between sleep quality and physical performance among a group of UK community-dwelling older adults, according to sex.

Methods: Sleep quality was assessed using the Pittsburgh Sleep Quality Index. Physical performance was assessed using a short physical performance battery (SPPB), a timed up-and-go, and a hand-grip strength test.

Results: Of 591 eligible study members, 401 completed the Pittsburgh Sleep Quality Index. In regression analyses, men who reported poor sleep quality were significantly more likely to have a poor SPPB score, even after adjustment for confounding factors (OR = 2.54, 95% CI 1.10–5.89, $P = .03$). The direction of the relationship was reversed among women, where those who reported poor sleep were less likely to have a low SPPB score (OR = 0.36, 95% CI 0.15–0.85, $P = .02$). Poor sleep quality was associated with poorer hand-grip strength among women (regression coefficient = -0.34 z score, 95% CI -0.64 , -0.04 , $P = .03$), but this relationship was not observed among men (regression coefficient = 0.28 z score, 95% CI -0.01 , 0.57 , $P = .06$).

Conclusion: We found evidence of an association between poor sleep quality and poorer physical performance in older adults, though there appear to be important sex differences.

© 2020 National Sleep Foundation. Published by Elsevier Inc. All rights reserved.

Introduction

Physical function declines with advancing age which can fundamentally affect health and well-being. When physical function becomes poor, it can increase the risk of falls and fractures,¹ and reduce an individual's ability to carry out everyday activities.² This, in turn, leads to a loss of autonomy and independence, and increases the costs of care and caregiver burden.³ Physical performance measures such as hand-grip strength and balance are primary indicators of overall physical function, and are independently predictive of future morbidity and mortality.^{4–6}

Sleep quality is also known to decline in later life.⁷ In the United Kingdom (UK), around a third of older adults report disturbed sleep.⁸ However, poor sleep quality among older people is not necessarily due to ageing alone. Multiple factors that are associated with the ageing process can affect sleep quality, for example, changes in physical or mental health, or changes in social engagement, lifestyle, and environment.⁹ Sleep quality is a multidimensional construct including sleep latency, wake after sleep onset, number and frequency of night awakenings, as well as subjective reports of feeling well-rested and refreshed upon awakening. Poor sleep quality in older people is associated with poorer quality of life,¹⁰ increased morbidity,¹¹ and increased mortality.¹²

As both physical performance and sleep quality decline with age, an association between them is plausible. Sleep may interact with

*Corresponding author.

E-mail address: h.denison@massey.ac.nz (H.J. Denison).

muscle mass and strength through metabolic, hormonal, and immunological factors,¹³ which may in turn have an effect on physical performance. For example, sleep deficit downregulates the hormones insulin-like growth factor 1 and testosterone which play central roles in protein synthesis pathways and thus maintenance of muscle mass.^{13,14} Several studies have demonstrated an association between poor physical performance and poor sleep, though most have been conducted in younger adults,¹⁵ athletes,¹⁶ or in clinical populations of older adults, such as those with osteoarthritis or cancer.^{17,18} Others have focused on the association between physical performance and sleep duration or specific sleep disorders such as sleep-disordered breathing, rather than sleep quality.^{19,20} Two particularly noteworthy studies are those by Goldman et al.²¹ and Dam et al.²² which both involved around 2800 participants in the United States; the first was among women only and the second among men only. These studies used objective measures of sleep (wrist actigraphy and polysomnography) and included physical performance measures such as walking speed, chair stands, and hand-grip strength. Both studies concluded that poorer sleep measures were associated with worse physical performance.

Few studies have explored sex differences in relation to sleep quality and physical performance outcomes,^{13,23,24} thus there is a limited understanding. To our knowledge, no studies that have assessed sex differences in the association have been conducted among UK populations. In fact, much of the research into the association of sleep and physical performance has been completed in the United States, China, and Japan, with fewer studies based on European populations. One study of sleep duration and physical performance has been conducted in the UK, and this was among older adults.²⁵ It showed that long sleep duration (>9 hours) was associated with poorer hand-grip strength and very slow walking speed at baseline, however differences between men and women were not explored. If interventions to improve physical performance or sleep are to be designed and targeted appropriately, it should not be assumed that the relationship will be identical across countries and/or ethnicities, or sex. We therefore sought to examine the potential association between sleep quality and physical performance characteristics in a large UK cohort of older adults and to assess differences in the association according to sex.

Methods

Study design and participants

Participants were recruited from the established Hertfordshire Cohort Study (HCS).²⁶ The HCS is a prospective, longitudinal, population-based study of the lifecourse origins of disease among people born in Hertfordshire, UK, between 1931 and 1939. In 2011–2012, participants were contacted to ask if they would be interested in taking part in a follow-up study. Those that agreed were visited at home by a trained research nurse, who administered a questionnaire and conducted a clinical examination. The participants were then given a second questionnaire which included a sleep assessment tool to complete in their own time and return to the study team in a prepaid envelope. The majority (73%) of the participants completed the second questionnaire on the same day as the nurse-assessment. The median (interquartile range [IQR]) time in days between the nurse-assessment and completing the second questionnaire was 0 (0–1), and the range was 0–40 days.

Sleep assessment

Sleep quality was assessed using the validated and reliable Pittsburgh Sleep Quality Index (PSQI).²⁷ The PSQI is comprised of 7 sub-components: (1) subjective sleep quality; (2) sleep latency (time

taken to initiate sleep); (3) sleep duration; (4) sleep efficiency (the ratio of total sleep time to time in bed); (5) sleep disturbance; (6) use of sleep-promoting medication (prescribed or over-the-counter); and (7) daytime dysfunction. Participants provide subjective sleep estimates based on the previous month. The answers were used to generate scores for each of the 7 sub-components, which ranged 0–3. A global PSQI score was generated by summing the scores of the sub-components ranging 0–21 with higher scores indicating poorer sleep quality. A score of more than 5 is indicative of poor sleep.²⁷ This cut-off has been shown to have high specificity and sensitivity for distinguishing poor sleepers and healthy controls.²⁷

Physical performance assessment

Physical performance was assessed using the short physical performance battery (SPPB),²⁸ a timed up-and-go (TUG) test and a hand-grip strength test. The SPPB is made up of a walking speed test, a chair rise test, and a standing balance test. To test walking speed, participants were asked to walk 8 feet (ft) at their customary pace while being timed using a stopwatch. The chair rise test involved participants moving from a sitting position (in a straight-backed chair) to a fully upright standing position 5 times as quickly as possible with their arms crossed across their chest while being timed. The standing balance test involved a semitandem stand where participants placed one foot in front of the other such that the big toe of one foot was touching the side of the heel of the other. If participants could not hold the semitandem stand for 10 seconds, they did a side-by-side stand (standing with the feet side-by-side). If they could hold the semitandem stand for 10 seconds, they also attempted a full tandem stand where participants placed one foot in front of the other (touching heel-to-toe) and held this position for as long as they could up to 10 seconds. A binary variable with a cut-off of 5 seconds was calculated for the full tandem stand, as it could not be analyzed continuously due to being censored at 10 seconds. Those who did not perform the full tandem stand because they could not hold the semitandem stand were recorded as 0 seconds and included in the tandem stand <5 seconds group for analysis. Performance on these tests was measured and reported individually, and also used to calculate a SPPB score.

The SPPB score was calculated as follows. For the walking speed and chair rise tests, those who could not complete the test were given a score of 0. The rest of the participants' times were divided into quartiles and given a score of 1–4, slowest to fastest quartile. For the balance test, if the participant maintained balance in the full tandem stand for 10 seconds, they were given a score of 4; if they obtained a time ≥ 3 and <10 seconds, they scored 3; if they obtained a time of <3 seconds but were able to maintain a semitandem stand, they scored 2; if they could not do the semitandem stand but could do the side-by-side stand, they scored 1; and if they could not do either the semitandem or the side-by-side stand, they scored 0. The scores were summed for the 3 measures, with a maximum of 12 and a minimum of 0. In keeping with previous work, those with a score equal to or lower than 9 were designated as having low physical performance.^{29,30}

Two other physical performance measures were used: a TUG test and a hand-grip strength test. For the TUG test, participants began seated in a chair with arms and were asked to stand, walk at their customary pace to a marked line at 3 m, then turn and walk back to sit in the chair while being timed.³¹ Hand-grip strength was measured using a Jamar hand-held isokinetic dynamometer while the participant sat in a chair with their wrist just over the end of the arm of the chair in a neutral position.³² The participants were given 3 attempts on each hand and the highest value measured in kilograms (kg) was used in analyses.

Covariates

Weight was measured to the nearest 0.1 kg using a calibrated scale, and height was measured to the nearest 0.001 m using a portable stadiometer. Body mass index (BMI) was calculated using the equation kg/m^2 . The nurse-administered questionnaire included items about demographics, general health, activity time (frequency and duration spent walking outside, cycling, gardening, doing sports, and carrying out household tasks in the last 2 weeks, converted into average minutes per day for analysis), and lifestyle (smoking and alcohol consumption). Alcohol consumption in units per week was calculated by converting the usual amount consumed of different alcoholic drinks (beer, wine, spirits/liqueurs) into units and multiplying this by the frequency of consumption of each type of alcoholic drink. As per previous research among this cohort,²⁶ social class was coded on the basis of current or most recent full-time occupation for men and never-married women, and on the basis of the husband's occupation for ever-married women, and categorized into 2 groups: classes I-IIIIM (nonmanual) and IIIM-V (manual) of the Standard Occupational Classification 1990 classification of social class. Participants were asked if they had any of the following comorbidities at the time of interview, and the total number was summed: high blood pressure, diabetes, respiratory disease, rheumatoid arthritis, multiple sclerosis, thyroid disease, vitiligo, depression, Parkinson's disease, heart disease, peripheral arterial disease, stroke, osteoporosis, and cancer. Participants were also asked if they regularly took any medications, and the number of these was summed.

Statistical analysis

Demographic, physical performance, and sleep characteristics of study participants were described using frequencies and percentages for dichotomous or categorical variables, and means and standard deviations or medians and IQR as appropriate for continuous variables.

The association of poor sleep quality (PSQI >5) with physical performance was assessed using logistic regression for dichotomous outcome variables (SPPB score and tandem stand), and linear regression for continuous outcome variables (8 ft walk, chair rises, TUG, and hand-grip strength). Chair rise time, TUG time, 8 ft walk, and hand-grip strength were transformed to Fisher-Yates *z* scores using the Fisher-Yates rank-based inverse normal transformation to normalize the data. We conducted all analyses for men and women separately due to significant interactions of 2 physical performance measures with gender (low overall physical performance [$P = .005$] and hand-grip strength [$P = .007$]), and our stated aim of exploring gender differences. Two models of adjustment were used. The first model adjusted for age, smoker status, alcohol consumption, social class, BMI, and activity time, as these were all significantly associated with at least one of the physical performance outcomes. The second model also adjusted for the number of comorbidities and the number of medications to assess whether these variables had any influence on the relationship between sleep quality and physical performance. These covariates, aside from social class and number of medications, have typically been included in models in similar studies.^{13,24} The analyses were repeated for each of the 7 subcomponents of the PSQI. A sensitivity analysis was conducted excluding participants who were taking any medications which may affect their sleep (including statins, calcium-channel blockers, beta-blockers, and codeine containing drugs). Cases were omitted from analyses where data were missing for variables included in the model (available case analysis).

The analyses were conducted with Stata version 14, and a *P* value of <.05 was considered statistically significant.

Ethical approval and consent

Ethical approval was granted by the Hertfordshire Research Ethics Committee, reference number 10/H0311/59, and all participants provided written, informed consent.

Results

A total of 591 HCS participants were eligible to participate, of whom 443 (75%) provided written informed consent to participate in the study. Of these, 406 (91%) returned the self-completed questionnaire. The PSQI was completed at least in part by 401 participants. A global PSQI score was calculated for those who had completed all of the PSQI subcomponents ($n = 373$). Where data were missing for one or more subcomponents ($N = 28$), a global PSQI score was not generated and the individual was excluded from analyses relating to global scores. However, these individuals were included in the analyses of subcomponents for the sections they had completed. Participant characteristics were compared between those with all the PSQI data, those with just some of the PSQI data and those with none of the PSQI data. There were no significant differences except that those with no PSQI data were more likely to be a current smoker and more likely to have poor tandem stand times (data not shown).

Table 1 shows the demographic characteristics of the 401 participants included in analyses. The mean age of men was 75.5 years and the mean age of women was 75.8 years. The mean BMI for men was 27.7 kg/m^2 and for women was 28.6 kg/m^2 . There were no statistically significant differences between men and women with respect to age, BMI, social class, the number of medications, or the number of comorbidities. Men had a higher alcohol consumption (median [IQR] 6.5 [1.0-14.0] units per week) compared to women (median [IQR] 0.5 [0.0-4.0] units per week), $P < .01$. Men were also more likely to have smoked than women (64.5% of women had never smoked compared to 40.3% of men), $P < .01$.

The physical performance and sleep characteristics of the study participants are shown in Table 2. The median SPPB score calculated from 3 of the tests (8 ft walk, chair rises, and tandem stands) was 9 for men, and 8 for women. With regards to sleep assessment, the median PSQI score was 4 for men and 5 for women, with 37% of men and 45% of women scoring more than 5, which is indicative of poor sleep.²⁷

The association of poor sleep quality (PSQI >5) with physical performance is reported in Table 3. Among men, poor sleep was associated with having a low overall SPPB score after adjustment for age, smoking, alcohol, social class, BMI, and activity time (OR = 2.60, 95% CI 1.15-5.89, $P = .02$). This association remained after additional adjustment for the number of comorbidities and the number of medications (OR = 2.54, 95% CI 1.10-5.89, $P = .03$). Subcomponent analyses indicated that in particular, more sleep disturbance was associated with a higher likelihood of a poor SPPB score for men (Table A, Appendix 1). The direction of the relationship was reversed among women, where poor sleep was associated with a lower likelihood of a poor SPPB score, though this association was only statistically significant when the number of comorbidities and medications were included in the adjustments (OR = 0.36, 95% CI 0.15-0.85, $P = .02$). Contrary to men, subcomponent analyses indicated that sleep duration and habitual sleep efficiency were associated with SPPB among women, where shorter duration and worse sleep efficiency were associated with better SPPB (Table A, Appendix 1). Overall poor sleep quality (PSQI >5) was not associated with 8 ft walk time, chair rise test, TUG time, or tandem stand performance among men or women. Among women, poor sleep was associated with a lower maximum hand-grip strength (regression coefficient = -0.37 *z* score, 95% CI $-0.66, -0.09$, $P = .01$); this remained significant after additional adjustment for the number of comorbidities and medications

Table 1
Demographic characteristics of the study participants

Characteristic	Men		Women		P value ^a
	Total n	Value	Total n	Value	
Age (years), mean (SD)	201	75.5 (2.5)	200	75.8 (2.6)	.29
BMI (kg/m ²), mean (SD)	200	27.7 (3.7)	197	28.6 (5.1)	.05
Social class, n (%)	191		200		.68
I-IIIIM		82 (42.9)		90 (45.0)	
IIIIM-V		109 (57.1)		110 (55.0)	
Activity time in last 2 weeks (min/day), median (IQR) ^b	187	181 (105-270)	191	200 (135-284)	.10
Alcohol consumption (units per week), median (IQR)	201	6.5 (1.0-14.0)	200	0.5 (0.0-4.0)	<.01*
Smoker status, n (%)	201		200		<.01*
Never		81 (40.3)		129 (64.5)	
Ex		112 (55.7)		67 (33.5)	
Current		8 (4.0)		4 (2.0)	
Number of medications, median (IQR)	201	5 (2-6)	200	4 (2-7)	.90
Number of comorbidities, n (%) ^c	201		200		.15
0		38 (18.9)		49 (24.5)	
1		74 (36.8)		56 (28.0)	
2		55 (27.4)		48 (24.0)	
3		19 (9.5)		23 (11.5)	
4 or more		15 (7.5)		24 (12.0)	

* Significant at the 5% level.

^aP value for difference between men and women.^bActivity time based on average minutes per day spent on walking outside, cycling, gardening, sports and household tasks in the last 2 weeks.^cNumber of comorbidities out of high blood pressure, diabetes, lung disease, rheumatoid arthritis, multiple sclerosis, thyroid disease, vitiligo, depression, Parkinson's disease, heart disease, peripheral arterial disease, stroke, osteoporosis, and cancer.

BMI, body mass index; IQR, interquartile range; n, number; SD, standard deviation.

(regression coefficient = -0.34 z score, 95% CI -0.64 , -0.04 , $P = .03$). The PSQI subcomponents subjective sleep quality and daytime dysfunction were associated with hand-grip strength after adjustment for all covariates (Table F, Appendix 1). No associations between hand-grip strength and poor overall sleep quality (PSQI >5) or any PSQI subcomponents were observed in men.

A sensitivity analysis was performed to exclude people who were on any medication which may affect their sleep. The pattern of the associations between poor sleep and physical performance was generally the same among this subgroup (data not shown). However, as the majority of people (70%) were taking medications that may affect their sleep, the sample size of people not taking sleep-affecting medication was very small (51 men and 69 women) meaning the analyses were underpowered to detect statistical significance or produce meaningful results.

Discussion

This study found evidence of a cross-sectional association between poor sleep quality and physical performance in men, where

men who reported poor sleep were significantly more likely to have a low overall SPPB score. This association was reversed for women, where women who reported poor sleep were less likely to have a low overall SPPB score. Interestingly, poor sleep quality was associated with poorer hand-grip strength among women, but this relationship was not observed among men.

It is interesting that the individual components of the SPPB (walking speed, chair rises, and standing balance) were not statistically significantly associated with sleep quality in either men or women, but that the composite SPPB score was. This finding suggests that overall physical functioning is related to sleep quality, rather than individual skills such as balance.

These results are similar to those from a cross-sectional study of older adults in Japan²⁴ where men who had poor sleep quality (measured by PSQI) had poorer physical performance than men with good sleep quality, but the same was not found among women. Conversely, Koh et al.²³ found that sleep quality (measured by PSQI) was associated with physical performance (SPPB score) among women and not men in their study of Korean older adults. Our finding that poor sleep quality was associated with poorer hand-grip strength in

Table 2
Physical performance and sleep characteristics of the study participants

Characteristic	Men		Women		P value ^a
	Total n	Value	Total n	Value	
SPPB score, median (IQR)	187	9 (7-11)	180	8 (7-10)	.01*
8 feet walk test (seconds), median (IQR)	191	3.0 (2.8-3.6)	190	3.2 (2.9-4.0)	<.01*
Chair rises (seconds), median (IQR)	178	15.7 (13.6-18.7)	171	16.7 (13.7-19.8)	.13
Tandem stand <5 s, n (%)	195	21 (10.8)	192	28 (14.6)	.26
Timed up-and-go (TUG) test (seconds), median (IQR)	188	11.3 (10.0-13.0)	187	11.9 (10.0-14.0)	.16
Maximum hand-grip strength (kg), median (IQR)	200	36.5 (32.0-41.0)	200	21.0 (18.0-26.0)	<.01*
PSQI score, median (IQR)	187	4 (2-7)	186	5 (3-9)	.02*
PSQI score >5, n (%)	187	69 (36.9)	186	84 (45.2)	.11

* Significant at the 5% level.

^aP value for difference between men and women.

IQR, interquartile range; n, number; PSQI, Pittsburgh Sleep Quality Index; SPPB, short physical performance battery.

Table 3
The association of poor sleep quality (PSQI >5) with physical performance

	Men				Women				
	Adjusted ^a		Adjusted ^b		Adjusted ^a		Adjusted ^b		
	n	Odds ratio (95% CI)	P value	n	Odds ratio (95% CI)	P value	n	Odds ratio (95% CI)	P value
Low SPPB score (≤9)	153	2.60 (1.15, 5.89)	.02*	153	2.54 (1.10, 5.89)	.03*	157	0.50 (0.23, 1.09)	.08
Poor tandem stand time (<5 s)	162	0.82 (0.28, 2.38)	.71	162	0.82 (0.27, 2.52)	.73	165	1.10 (0.42, 2.92)	.84
	n	Regression coefficient (95% CI)	P value	n	Regression coefficient (95% CI)	P value	n	Regression coefficient (95% CI)	P value
8 feet walk time ^c	155	0.11 (-0.21, 0.42)	.50	155	0.07 (-0.25, 0.39)	.66	165	-0.02 (-0.28, 0.25)	.91
Chair rise time ^c	147	0.27 (-0.04, 0.59)	.08	147	0.28 (-0.03, 0.58)	.08	148	-0.19 (-0.49, 0.11)	.22
Timed up-and-go (TUG) time ^c	153	0.23 (-0.05, 0.51)	.11	153	0.18 (-0.10, 0.47)	.21	162	-0.04 (-0.31, 0.23)	.78
Max hand-grip strength ^c	163	0.24 (-0.04, 0.52)	.09	163	0.28 (-0.01, 0.57)	.06	174	-0.37 (-0.66, -0.09)	.01*
	n	Regression coefficient (95% CI)	P value	n	Regression coefficient (95% CI)	P value	n	Regression coefficient (95% CI)	P value
8 feet walk time ^c	155	0.11 (-0.21, 0.42)	.50	155	0.07 (-0.25, 0.39)	.66	165	-0.15 (-0.41, 0.12)	.28
Chair rise time ^c	147	0.27 (-0.04, 0.59)	.08	147	0.28 (-0.03, 0.58)	.08	148	-0.30 (-0.60, 0.01)	.06
Timed up-and-go (TUG) time ^c	153	0.23 (-0.05, 0.51)	.11	153	0.18 (-0.10, 0.47)	.21	162	-0.13 (-0.41, 0.14)	.34
Max hand-grip strength ^c	163	0.24 (-0.04, 0.52)	.09	163	0.28 (-0.01, 0.57)	.06	174	-0.34 (-0.64, -0.04)	.03*

^a Fisher Yates z score.

* Significant at the 5% level.

^b Adjusted for age, smoking, alcohol, social class, BMI, and activity time.

^c Adjusted for age, smoking, alcohol, social class, BMI, activity time, number of comorbidities, and number of medications.

SPPB, short physical performance battery.

women but not men is similar to findings of a study of older people in Germany,¹³ where women with poor sleep had reduced hand-grip strength but no association was seen among men.

While these previous studies have found associations between sleep and physical performance in one sex but not the other, our study is the first to find a difference in directionality between the sexes for the association between sleep and overall SPPB score. There are a number of possible explanations for these findings. The difference could be due to the men and women in the study subjectively assessing their sleep differently so that the pattern of association with physical performance appears to be different.³³ Changes in sex hormones with age which affect sleep and physiology may also have an impact on the differences observed between men and women. Alternatively, this result could be a spurious association due to the number of statistical tests conducted. Further research is needed to determine whether this is a robust finding or due to chance.

That women with poor sleep quality were less likely to have a low overall SPPB score but more likely to have a poor hand-grip strength score appears discrepant. One hypothesis is that this inconsistency may be related to the physiological areas involved in these tasks (the SPPB primarily assesses lower extremity functioning whereas the hand-grip strength test is focused on the upper extremity of the body) if the mechanism by which sleep and physical performance are associated acts differently on different parts of the body. The authors are not aware of any studies investigating the physiological mechanisms behind the association between sleep quality and physical performance.

To our knowledge, this is the first study to examine the association between sleep quality and physical performance among community-dwelling older adults in the UK. Previous research has predominantly been conducted in the United States, China, and Japan. One of the strengths of this study is the sampled population. The individuals recruited were a nonselective community-dwelling population who were born in Hertfordshire, UK. HCS characteristic and mortality patterns have previously been demonstrated to be similar to those of the nationally representative Health Survey for England.²⁶ However, it should be noted that the sample included in this analysis does not represent the whole of the HCS cohort. Additionally, the cohort is predominantly White so results may not be generalizable to other ethnic groups. As with all cohort studies there is also the possibility of healthy responder bias, where the more unwell members of the population contacted are less likely to participate due to their ill-health.

A limitation of this study is that it does not include a measure of sleep-disordered breathing. Sleep-disordered breathing causes hypoxia which in turn may lead to muscle wasting and weakness.³⁴ The average BMI of both the men and women in this group was in the overweight range, so it is possible that obstructive sleep apnea (the most common form of sleep-disordered breathing) was contributing to poor sleep quality and physical decline. To address this, models were adjusted for BMI in our analyses.

The sleep quality data in the current study were self-reported, and therefore may have been affected by biases including social desirability or selective recall bias. This may be especially relevant for the sub-components, which are based on single items. While multnight laboratory polysomnography is the most accurate method of measuring sleep, it is extremely resource intensive for a large sample. Self-reported sleep data are commonly used for large studies, and the PSQI is a well-utilized tool for assessment of sleep quality. While the PSQI cannot be used to screen for polysomnographic sleep abnormalities,³⁵ it has been shown to be a useful, valid and reliable tool for the assessment of habitual sleep quality and is often independently associated with health outcomes.^{36–38}

A further limitation of this study is that, in the analysis of the association between sleep and physical performance, some of the

participants had missing data for physical performance measures or adjustment variables. To maximize power, we used available case analysis where each individual model included all participants who had complete data for that model (sample sizes presented in Table 3). This means that the samples differ across analyses, which could have led to some bias in results and reduces generalizability.

While we have shown that physical performance and sleep could be related among older men and women, the cross-sectional nature of the study prevents the assessment of whether this is a causal relationship or of the longitudinal nature of the relationship. However, a few studies have assessed the relationship between sleep and physical performance longitudinally.^{39,40} A prospective study of older women reported that objectively measured poorer sleep quality predicts substantial decline in hand-grip strength over 5 years, but not decline in gait speed.³⁹ Similar findings were reported for the association between baseline sleep quality and activities of daily living functioning at 6-month follow-up among residents of assisted living facilities.⁴⁰ While there have been several randomized controlled studies measuring the effects of sleep on physical performance among younger adults and athletes,^{15,41} the authors are not aware of any such intervention studies among older adults.

In conclusion, we found evidence of an association between poor sleep quality and physical performance in UK community-dwelling older adults, as has been reported by studies in other countries. These results also demonstrate important sex differences in the relationship between sleep quality and physical performance. Further prospective longitudinal studies among community-dwelling older adults to confirm these cross-sectional findings are warranted.

Funding

The Hertfordshire Cohort Study was supported by the Medical Research Council of Great Britain; Arthritis Research UK; the Porticus Foundation, and the International Osteoporosis Foundation. The work herein was also supported by the NIHR Nutrition BRC, University of Southampton and the NIHR Musculoskeletal BRU, University of Oxford. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests

The authors have no competing interests to declare.

Acknowledgments

The research team would like to thank the individuals who generously shared their time to participate in this study. We would also like to thank the HCS Research Staff including the fieldworkers who worked on the study, and Vanessa Cox and Ken Cox who managed the data.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:10.1016/j.sleh.2020.10.002.

References

1. VanSwearingen JM, Paschal KA, Bonino P, Chen TW. Assessing recurrent fall risk of community-dwelling, frail older veterans using specific tests of mobility and the physical performance test of function. *J Gerontol A Biol Sci Med Sci*. 1998;53:M457–M464.
2. Brach JS, VanSwearingen JM. Physical impairment and disability: relationship to performance of activities of daily living in community-dwelling older men. *Phys Ther*. 2002;82:752–761.
3. Cole CS, Richards KC, Beck CC, et al. Relationships among disordered sleep and cognitive and functional status in nursing home residents. *Res Gerontol Nurs*. 2009;2:183–191.
4. Simmonds SJ, Syddall HE, Westbury LD, Dodds RM, Cooper C, Aihie Sayer A. Grip strength among community-dwelling older people predicts hospital admission during the following decade. *Age Ageing*. 2015;44:954–959.
5. Granic A, Davies K, Jagger C, Dodds RM, Kirkwood TBL, Sayer AA. Initial level and rate of change in grip strength predict all-cause mortality in very old adults. *Age Ageing*. 2017;46:970–976.
6. Keevil VL, Luben R, Hayat S, Sayer AA, Wareham NJ, Khaw KT. Physical capability predicts mortality in late mid-life as well as in old age: findings from a large British cohort study. *Arch Gerontol Geriatr*. 2018;74:77–82.
7. Crowley K. Sleep and sleep disorders in older adults. *Neuropsychol Rev*. 2011;21:41–53.
8. Jackowska M, Kumari M, Steptoe A. Sleep and biomarkers in the English Longitudinal Study of Ageing: associations with C-reactive protein, fibrinogen, dehydroepiandrosterone sulfate and hemoglobin. *Psychoneuroendocrinology*. 2013;38:1484–1493.
9. Li J, Vitiello MV, Gooneratne NS. Sleep in normal aging. *Sleep Med Clin*. 2018;13:1–11.
10. Lee M, Choh AC, Demerath EW, et al. Sleep disturbance in relation to health-related quality of life in adults: the Fels Longitudinal Study. *J Nutr Health Aging*. 2009;13:576–583.
11. von Ruesten A, Weikert C, Fietze I, Boeing H. Association of sleep duration with chronic diseases in the European Prospective Investigation into Cancer and Nutrition (EPIC)-Potsdam study. *PLoS One*. 2012;7:e30972.
12. Cappuccio FP, D'Elia L, Strazzullo P, Miller MA. Sleep duration and all-cause mortality: a systematic review and meta-analysis of prospective studies. *Sleep*. 2010;33:585–592.
13. Buchmann N, Spiira D, Norman K, Demuth I, Eckardt R, Steinhagen-Thiessen E. Sleep, muscle mass and muscle function in older people. *Dtsch Arztebl Int*. 2016;113:253–260.
14. Dattilo M, Antunes HK, Medeiros A, et al. Sleep and muscle recovery: endocrinological and molecular basis for a new and promising hypothesis. *Med Hypotheses*. 2011;77:220–222.
15. Patrick Y, Lee A, Raha O, et al. Effects of sleep deprivation on cognitive and physical performance in university students. *Sleep Biol Rhythms*. 2017;15:217–225.
16. Fullagar HH, Skorski S, Duffield R, Hammes D, Coutts AJ, Meyer T. Sleep and athletic performance: the effects of sleep loss on exercise performance, and physiological and cognitive responses to exercise. *Sports Med*. 2015;45:161–186.
17. Wilcox S, Brenes GA, Levine D, Sevick MA, Shumaker SA, Craven T. Factors related to sleep disturbance in older adults experiencing knee pain or knee pain with radiographic evidence of knee osteoarthritis. *J Am Geriatr Soc*. 2000;48:1241–1251.
18. Loh KP, Pandya C, Zittel J, et al. Associations of sleep disturbance with physical function and cognition in older adults with cancer. *Support Care Cancer*. 2017;25:3161–3169.
19. Chen HC, Hsu NW, Chou P. The association between sleep duration and hand grip strength in community-dwelling older adults: the Yilan Study, Taiwan. *Sleep*. 2017;40.
20. Suri SV, Batterham AM, Ells L, Danjoux G, Atkinson G. Cross-sectional association between walking pace and sleep-disordered breathing. *Int J Sports Med*. 2015;36:843–847.
21. Goldman SE, Stone KL, Ancoli-Israel S, et al. Poor sleep is associated with poorer physical performance and greater functional limitations in older women. *Sleep*. 2007;30:1317–1324.
22. Dam TT, Ewing S, Ancoli-Israel S, Ensrud K, Redline S, Stone K. Association between sleep and physical function in older men: the osteoporotic fractures in men sleep study. *J Am Geriatr Soc*. 2008;56:1665–1673.
23. Koh ES, Jang SN, Paik NJ, Kim KW, Lim JY. Age and gender patterns in associations between lifestyle factors and physical performance in older Korean adults. *Arch Gerontol Geriatr*. 2014;59:338–345.
24. Malinowska KB, Ikezoe T, Ichihashi N, et al. Self-reported quality of sleep is associated with physical strength among community-dwelling young-old adults. *Geriatr Gerontol Int*. 2017;17:1808–1813.
25. Morgan K, Harteescu I. Sleep duration and all-cause mortality: links to physical activity and prefrailty in a 27-year follow up of older adults in the UK. *Sleep Med*. 2019;54:231–237.
26. Syddall HE, Aihie Sayer A, Dennison EM, Martin HJ, Barker DJ, Cooper C. Cohort profile: the hertfordshire cohort study. *Int J Epidemiol*. 2005;34:1234–1242.
27. Buysse DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res*. 1989;28:193–213.
28. Guralnik JM, Simonsick EM, Ferrucci L, et al. A short physical performance battery assessing lower extremity function: association with self-reported disability and prediction of mortality and nursing home admission. *J Gerontol*. 1994;49:M85–M94.
29. Abbatecola AM, Cherubini A, Guralnik JM, et al. Plasma polyunsaturated fatty acids and age-related physical performance decline. *Rejuvenation Res*. 2009;12:25–32.
30. da Camara SM, Alvarado BE, Guralnik JM, Guerra RO, Maciel AC. Using the short physical performance battery to screen for frailty in young-old adults with distinct socioeconomic conditions. *Geriatr Gerontol Int*. 2013;13:421–428.
31. Podsiadlo D, Richardson S. The timed "Up & Go": a test of basic functional mobility for frail elderly persons. *J Am Geriatr Soc*. 1991;39:142–148.

32. Roberts HC, Denison HJ, Martin HJ, et al. A review of the measurement of grip strength in clinical and epidemiological studies: towards a standardised approach. *Age Ageing*. 2011;40:423–429.
33. van den Berg JF, Miedema HME, Tulen JHM, Hofman A, Neven AK, Tiemeier H. Sex differences in subjective and actigraphic sleep measures: a population-based study of elderly persons. *Sleep*. 2009;32:1367–1375.
34. Wüst RCI, Degens H. Factors contributing to muscle wasting and dysfunction in COPD patients. *Int J Chronic Obstructive Pulm Dis*. 2007;2:289–300.
35. Buysse DJ, Hall ML, Strollo PJ, et al. Relationships between the Pittsburgh Sleep Quality Index (PSQI), Epworth Sleepiness Scale (ESS), and clinical/polysomnographic measures in a community sample. *J Clin Sleep Med*. 2008;4:563–571.
36. Beaudreau SA, Spira AP, Stewart A, et al. Validation of the Pittsburgh Sleep Quality Index and the Epworth Sleepiness Scale in older black and white women. *Sleep Med*. 2012;13:36–42.
37. Spira AP, Beaudreau SA, Stone KL, et al. Reliability and validity of the Pittsburgh Sleep Quality Index and the Epworth Sleepiness Scale in older men. *J Gerontol A Biol Sci Med Sci*. 2012;67:433–439.
38. Paine S-J, Harris R, Cormack D, Stanley J. Self-reported sleep complaints are associated with adverse health outcomes: cross-sectional analysis of the 2002/03 New Zealand Health Survey. *Ethnicity Health*. 2019;24:44–56.
39. Spira AP, Covinsky K, Rebok GW, et al. Poor sleep quality and functional decline in older women. *J Am Geriatr Soc*. 2012;60:1092–1098.
40. Martin JL, Fiorentino L, Jouldjian S, Josephson KR, Alessi CA. Sleep quality in residents of assisted living facilities: effect on quality of life, functional status, and depression. *J Am Geriatr Soc*. 2010;58:829–836.
41. Fullagar H, Skorski S, Duffield R, Meyer T. The effect of an acute sleep hygiene strategy following a late-night soccer match on recovery of players. *Chronobiol Int*. 2016;33:490–505.